

The Membership Relation in Fuzzy Operators

ZHANG Shiqiang^{[a],[b],*}, and LUO Yaling^{[a],[b]}

^[a] Department of Mathematics, College of Basic Medical Sciences, Chongqing Medical University, Chongqing, China.

^[b] Lab. of Forensic Medicine and Biomedicine Information, Chongqing Medical University, Chongqing, China.

* Corresponding author.

Address: Department of Mathematics, College of Basic Medical Sciences, & Lab. of Forensic Medicine and Biomedicine Information, Chongqing Medical University, Chongqing, 400016, China; E-Mail: math808@sohu.com

Supported by College of Basic Medical Sciences of Chongqing Medical University, Grant No. 2011. **Special thanks to** Science and Technology Committee of Yuzhong District (STCYD) of Chongqing as this paper is based on the Science and Technology Plan Projects “Research on Birth Defects Intervention Information System for ‘Health Chongqing’ ” (code No. 20110403) supported by STCYD.

Received: August 14, 2012/ Accepted: October 4, 2012/ Published: November 30, 2012

Abstract: Firstly the membership relation in fuzzy operators located outside Zadeh operators be discussed and given. Secondly the membership relation in fuzzy operators located within Zadeh operators be discussed and given.

Key words: Operator; Fuzzy operator; Membership relation

ZHANG, S., & LUO, Y. (2012). The Membership Relation in Fuzzy Operators. *Studies in Mathematical Sciences*, 5(2), 98–104. Available from <http://www.cscanada.net/index.php/sms/article/view/j.sms.1923845220120502.ZR0217> DOI: 10.3968/j.sms.1923845220120502.ZR0217

1. INTRODUCTION

We since Zadeh established fuzzy set theorem in 1965, he introduced a pair of fuzzy operators. They are named Zadeh operators [1]. There is a lot of discussion about the fuzzy operators [2–10], but there is little discussion membership relation in there

operators. This paper discusses membership relation in Zadeh operators and other generalized operators in common use.

2. THE MEMBERSHIP RELATION IN FUZZY OPERATORS LOCATED OUTSIDE ZADEH OPERATORS

In order to discuss the membership relation in fuzzy operators, it is necessary to introduce the concept of generalized operators [4]. Generalized operators are widely-used operators in fuzzy sets. The definition of Zadeh operators and the definitions of generalized operators in common use as well as their membership relation are as follows.

Definition 2.1 Zadeh operators ($\wedge, \vee$)

To any fuzzy set $A, B, C \in P(U)$, fuzzy set $C = A \cap B$ be called intersection of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$C(u) = (A \cap B)(u) = A(u) \wedge B(u);$$

Fuzzy set $C = A \cup B$ be called union of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$C(u) = (A \cup B)(u) = A(u) \vee B(u).$$

Definition 2.2 Probability operators ($\hat{\bullet}, \hat{+}$)

To any fuzzy set $A, B, C \in P(U)$, fuzzy set $C = A \hat{\bullet} B$ be called probability product of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$C(u) = (A \hat{\bullet} B)(u) = A(u)B(u);$$

Fuzzy set $C = A \hat{+} B$ be called probability sum of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$C(u) = (A \hat{+} B)(u) = A(u) + B(u) - A(u)B(u).$$

Definition 2.3 Boundary operators ($\otimes, \oplus$)

To any fuzzy set $A, B, C \in P(U)$, fuzzy set $C = A \otimes B$ be called boundary product of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$C(u) = (A \otimes B)(u) = \max[0, A(u) + B(u) - 1];$$

Fuzzy set $C = A \oplus B$ be called boundary sum of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$C(u) = (A \oplus B)(u) = \min[1, A(u) + B(u)].$$

Definition 2.4 Infinite operators ($\hat{\infty}, \hat{\infty}$)

To any fuzzy set $A, B, C \in P(U)$, fuzzy set $C = A \hat{\infty} B$ be called infinite product of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$\tilde{C}(u) = (\tilde{A} \hat{\infty} \tilde{B})(u) = \begin{cases} \tilde{A}(u), & \tilde{B}(u) = 1 \\ \tilde{B}(u), & \tilde{A}(u) = 1 \\ 0, & \text{other} \end{cases}$$

Fuzzy set $C = A \widetilde{\infty} B$ be called infinite sum of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$\tilde{C}(u) = (\tilde{A} \widetilde{\infty} \tilde{B})(u) = \begin{cases} \tilde{A}(u), & \tilde{B}(u) = 0 \\ \tilde{B}(u), & \tilde{A}(u) = 0 \\ 1, & \text{other} \end{cases}$$

The paper [7] has proved that Zadeh operations, probability operators and boundary operators as well as infinite operators have membership relation as follows:

$$\begin{aligned} (\tilde{A} \hat{\infty} \tilde{B})(u) &\leq (\tilde{A} \otimes \tilde{B})(u) \leq (\tilde{A} \bullet \tilde{B})(u) \leq (\tilde{A} \cap \tilde{B})(u) \\ &\leq (\tilde{A} \cup \tilde{B})(u) \leq (\tilde{A} \hat{+} \tilde{B})(u) \leq (\tilde{A} \oplus \tilde{B})(u) \leq (\tilde{A} \widetilde{\infty} \tilde{B})(u) \end{aligned}$$

Definition 2.5 Schweizer-Sklard operators $(\overset{\bullet}{ss}, \overset{+}{ss})$ [11]

To any fuzzy set $A, B, C \in P(U)$, $p \rightarrow (-\infty, +\infty)$, fuzzy set $C = A \overset{\bullet}{ss} B$ be called Schweizer-Sklard product of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$\tilde{A}(u) \overset{\bullet}{ss} \tilde{B}(u) = \begin{cases} 0, & \text{if } p > 0, \tilde{A}(u) = 0 \text{ or } \tilde{B}(u) = 0 \\ [\tilde{A}(u)^{-p} + \tilde{B}(u)^{-p} - 1]^{-1/p}, & \text{if } p > 0, \tilde{A}(u) \bullet \tilde{B}(u) \neq 0 \\ \tilde{A}(u) \bullet \tilde{B}(u), & \text{if } p = 0 \\ [\tilde{A}(u)^{-p} + \tilde{B}(u)^{-p} - 1]^{-1/p}, & \text{if } p < 0, \tilde{A}(u)^{-p} + \tilde{B}(u)^{-p} > 1 \\ 0, & \text{if } p < 0, \tilde{A}(u)^{-p} + \tilde{B}(u)^{-p} \leq 1 \end{cases}$$

Fuzzy set $C = A \overset{+}{ss} B$ be called Schweizer-Sklard sum of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$\tilde{A}(u) \overset{+}{ss} \tilde{B}(u) = \begin{cases} 1, & \text{if } p > 0, \tilde{A}(u) = 1 \text{ or } \tilde{B}(u) = 1 \\ 1 - \{[1 - \tilde{A}(u)]^{-p} + [1 - \tilde{B}(u)]^{-p} - 1\}^{-1/p}, & \text{if } p > 0, \tilde{A}(u) \bullet \tilde{B}(u) \neq 1 \\ \tilde{A}(u) + \tilde{B}(u) - \tilde{A}(u) \bullet \tilde{B}(u), & \text{if } p = 0 \\ 1 - \{[1 - \tilde{A}(u)]^{-p} + [1 - \tilde{B}(u)]^{-p} - 1\}^{-1/p}, & \text{if } p < 0, [1 - \tilde{A}(u)]^{-p} + [1 - \tilde{B}(u)]^{-p} > 1 \\ 0, & \text{if } p < 0, [1 - \tilde{A}(u)]^{-p} + [1 - \tilde{B}(u)]^{-p} \leq 1 \end{cases}$$

The paper [11] has proved that Schweizer-Sklard operators are monotone functions for variables $A(u)$ or $B(u)$, the paper [11] has proved that Schweizer-Sklard operators are monotone functions for the parameter p .

The paper [11] has proved that Schweizer-Sklard operations, and Zadeh operations as well as probability operators have membership relation as follows:

$$\begin{aligned}\tilde{A}(u) \dot{\circ} \tilde{B}(u) &\subseteq \tilde{A}(u) \overset{\bullet}{ss} \tilde{B}(u) \subseteq \tilde{A}(u) \wedge \tilde{B}(u) \\ &\subseteq \tilde{A}(u) \vee \tilde{B}(u) \subseteq \tilde{A}(u) \overset{+}{ss} \tilde{B}(u) \subseteq \tilde{A}(u) \hat{+} \tilde{B}(u)\end{aligned}$$

Definition 2.6 Dobois-Prade operators $(\overset{\bullet}{d}, \overset{+}{d})$ [13]

To any fuzzy set $A, B, C \in P(U)$, $\lambda \in [0, 1]$, fuzzy set $C = A \overset{\bullet}{d} B$ be called Dobois-Prade product of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$(\tilde{A} \overset{\bullet}{d} \tilde{B})(u) = \begin{cases} 0, & \text{if } \lambda = 0 \text{ and } \tilde{A}(u) = \tilde{B}(u) = 0 \\ \frac{\tilde{A}(u)\tilde{B}(u)}{\max\{\lambda, \tilde{A}(u), \tilde{B}(u)\}}, & \text{others} \end{cases}$$

Fuzzy set $C = A \overset{+}{d} B$ be called Dobois-Prade sum of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$(\tilde{A} \overset{+}{d} \tilde{B})(u) = \begin{cases} 1, & \text{if } \lambda = 0 \text{ and } \tilde{A}(u) = \tilde{B}(u) = 1 \\ \frac{\tilde{A}(u) + \tilde{B}(u) - \tilde{A}(u)\tilde{B}(u) - \min\{1 - \lambda, \tilde{A}(u), \tilde{B}(u)\}}{\max\{\lambda, 1 - \tilde{A}(u), 1 - \tilde{B}(u)\}}, & \text{others} \end{cases}$$

The paper [13] has proved that Dobois-Prade operators are monotone functions for the parameter λ and that Dobois-Prade operations, and Zadeh operations as well as probability operators have membership relation as follows:

$$(\tilde{A} \hat{\circ} \tilde{B})(u) \leq (\tilde{A} \overset{\bullet}{d} \tilde{B})(u) \leq (\tilde{A} \cap \tilde{B})(u) \leq (\tilde{A} \cup \tilde{B})(u) \leq (\tilde{A} \overset{+}{d} \tilde{B})(u)$$

Definition 2.7 Hamacher operators $(\overset{\bullet}{r}, \overset{+}{r})$ [9]

To any fuzzy set $A, B, C \in P(U)$, $r \in [0, +\infty)$, fuzzy set $C = A \overset{\bullet}{r} B$ be called Hamacher product of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$(\tilde{A} \overset{\bullet}{r} \tilde{B})(u) = \begin{cases} 0, & \text{if } r = 0 \text{ and } \tilde{A}(u) = \tilde{B}(u) = 0 \\ \frac{\tilde{A}(u)\tilde{B}(u)}{r + [1 - r][\tilde{A}(u) + \tilde{B}(u) - \tilde{A}(u)\tilde{B}(u)]}, & \text{others} \end{cases}$$

Fuzzy set $C = A \overset{+}{r} B$ be called Hamacher sum of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$(\tilde{A} \overset{+}{r} \tilde{B})(u) = \begin{cases} 1, & \text{if } r = 0, \text{ and } \tilde{A}(u) = \tilde{B}(u) = 1 \\ \frac{\tilde{A}(u) + \tilde{B}(u) + [r - 2]\tilde{A}(u)\tilde{B}(u)}{1 - \tilde{A}(u)\tilde{B}(u) + r\tilde{A}(u)\tilde{B}(u)}, & \text{others} \end{cases}$$

The paper [9] has proved that Hamacher operations are monotone functions for the parameter r and that Hamacher operations, and Zadeh operations as well as boundary operators have membership relation as follows:

$$(\tilde{A} \otimes \tilde{B})(u) \leq (\tilde{A} \overset{\bullet}{\tilde{r}} \tilde{B})(u) \leq (\tilde{A} \cap \tilde{B})(u) \leq (\tilde{A} \cup \tilde{B})(u) \leq (\tilde{A} \overset{+}{\tilde{r}} \tilde{B})(u) \leq (\tilde{A} \oplus \tilde{B})(u)$$

Definition 2.8 Yager operators $(\overset{\bullet}{\Upsilon}, \overset{+}{\Upsilon})$ [10]

To any fuzzy set $A, B, C \in P(U)$, $r \in (0, +\infty)$, fuzzy set $C = A \overset{\bullet}{\Upsilon} B$ be called Yager product of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$\tilde{C}(u) = (\tilde{A} \overset{\bullet}{\Upsilon} \tilde{B})(u) = 1 - 1 \wedge [(1 - \tilde{A}(u))^{\frac{1}{r}} + (1 - \tilde{B}(u))^{\frac{1}{r}}]^r$$

Fuzzy set $C = A \overset{+}{\Upsilon} B$ be called Yager sum of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$\tilde{C}(u) = (\tilde{A} \overset{+}{\Upsilon} \tilde{B})(u) = 1 \wedge [\tilde{A}(u)^{\frac{1}{r}} + \tilde{B}(u)^{\frac{1}{r}}]^r$$

The paper [10] has proved that Yager operations are monotone functions for the parameter r and that Yager operations, and Zadeh operations as well as boundary operators have membership relation as follows:

$$(\tilde{A} \otimes \tilde{B})(u) \leq (\tilde{A} \overset{\bullet}{\Upsilon} \tilde{B})(u) \leq (\tilde{A} \cap \tilde{B})(u) \leq (\tilde{A} \cup \tilde{B})(u) \leq (\tilde{A} \overset{+}{\Upsilon} \tilde{B})(u) \leq (\tilde{A} \oplus \tilde{B})(u)$$

3. THE MEMBERSHIP RELATION IN FUZZY OPERATORS LOCATED WITHIN ZADEH OPERATORS

The paper [12] introduce new class of fuzzy operators in inner of Zadeh operator. The definitions of the new class of continuous fuzzy operators located within Zadeh operators and their membership membership relation are as follows:

Definition 3.1 Zero operators $(\hat{0}, \tilde{0})$

To any fuzzy set $A, B, C \in P(U)$, fuzzy set $C = A \hat{0} B$ be called Zero product of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$\tilde{C}(u) = (\tilde{A} \hat{0} \tilde{B})(u) = \frac{\tilde{A}(u) + \tilde{B}(u)}{2}$$

Fuzzy set $C = A \tilde{0} B$ be called Zero sum of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$\tilde{C}(u) = (\tilde{A} \tilde{0} \tilde{B})(u) = \frac{\tilde{A}(u) + \tilde{B}(u)}{2}$$

The paper [12] has proved that Zero operations and Zadeh operations have membership relation as follows:

$$(\tilde{A} \cap \tilde{B})(u) < (\tilde{A} \hat{0} \tilde{B})(u) = (\tilde{A} \tilde{0} \tilde{B})(u) < (\tilde{A} \cup \tilde{B})(u)$$

Definition 3.2 Q operators $(\hat{Q}, \tilde{Q})$

To any fuzzy set $A, B, C \in P(U)$, fuzzy set $C = A\hat{Q}B$ be called Q product of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$\tilde{C}(u) = (\tilde{A}\hat{Q}\tilde{B})(u) = \frac{2q+1}{4q+1}[\tilde{A}(u) \wedge \tilde{B}(u)] + \frac{2q}{4q+1}[\tilde{A}(u) \vee \tilde{B}(u)]$$

Fuzzy set $C = A\tilde{Q}B$ be called Q sum of fuzzy set A and fuzzy set B if $\forall u \in U$, there is

$$\tilde{C}(u) = (\tilde{A}\tilde{Q}\tilde{B})(u) = \frac{2q}{4q+1}[\tilde{A}(u) \wedge \tilde{B}(u)] + \frac{2q+1}{4q+1}[\tilde{A}(u) \vee \tilde{B}(u)]$$

The Q operators are monotone functions for the parameter q [12]. The paper [12] has proved that Q operations, Zadeh operations as well as Zero operators have membership relation as follows:

$$(\tilde{A} \cap \tilde{B})(u) \leq (\tilde{A}\hat{Q}\tilde{B})(u) \leq (\tilde{A}\hat{O}\tilde{B})(u) = (\tilde{A}\tilde{O}\tilde{B})(u) \leq (\tilde{A}\tilde{Q}\tilde{B})(u) \leq (\tilde{A} \cup \tilde{B})(u)$$

4. CONCLUSIONS

First we have discussed and given the membership relation in fuzzy operators located outside Zadeh operators. Second we have discussed and given the membership relation in fuzzy operators located within Zadeh operators.

REFERENCES

- [1] Zadeh, L. A. (1965). Fuzzy sets. *Information and Control*, (8), 338-359.
- [2] LI, H. X., & WANG, P. Z. (1994). *Fuzzy mathematics* (pp. 128-129). Beijing: National Defence Industry Press.
- [3] CHENG, Y. Y. (1984). *Fuzzy mathematics* (pp. 30-36). Wuhan: Huazhong Institute of Technology Press.
- [4] Lou, C. Z. (1989). *Introductory fuzzy sets* (pp. 44-56). Beijing: Beijing Teachers University Press.
- [5] SU, Z. F. (1989). *Fuzzy mathematics and medical science*. Beijing: Scientific and Technical Documents Press.
- [6] HAN, L. Y., & WANG, P. Z. (1998). *Applied fuzzy mathematics* (pp. 148-154). Beijing: Capital Economy and Trade University Press.
- [7] ZHANG, Shiqiang (2008). Method of checking up concealed faults on the fuzzy controller. *Proc. of the China Association for Science and Technology*, 4(1), 59-63.
- [8] ZHANG, Shiqiang, ZHANG, Salan, & JIANG, Zheng (2010). The problem and countermeasures on construction of fuzzy synthetic decision model. *International Conference on Engineering and Business Management, Scientific Research Publishing*, 1259-1260.
- [9] ZHANG, Shiqiang (2007). A new specific property of Hamacher operators and its application. *Fuzzy Systems and Mathematics*, 21(5), 83-88.
- [10] ZHANG, Shiqiang, & ZHANG, Salan (2010). A new specific property of Yager operators and its application. *International Conference on Computer, Mechatronics, Control and Electronic Engineering, IEEE*, 1, 144-147.
- [11] WANG, Chunli, & ZHANG, Shiqiang (2011). A new specific property of Schweizer-Sklard operators. *Studies in Mathematical Sciences*, 3(1), 52-57.

- [12] ZHANG, Shiqiang, & ZHANG, Salan (2012a). Fuzzy zero operator, fuzzy Q operator and its properties. *The 2rd International Conference on Electrics and Electronics*, 805-812.
- [13] ZHANG, Shiqiang, & ZHANG, Salan (2012b). A new specific property of Dobois-Prade operators. *The 2rd International Conference on Electrics and Electronics*, 813-820.